

*Liste des synonymes.*

<i>arcuata</i>	1910	(Gahvagni)	=	<i>dentata</i>	1915	(Bryk)	C 2
<i>amplusdecora</i>	1929	(Eisner)	=	<i>lunigera</i>	1911	(Turati)	
			=	<i>rubrocatenata</i>	1929	(Stauder)	A 2
<i>chryseis</i>	1911	(Turati)	=	<i>theoides</i>	1912	(Schwawerda)	D
<i>depupillata</i>	1918	(Turati)	=	<i>expupillata</i>	1911	(Rocci)	B 1
<i>fasciata</i>	1906	(Stichel)	=	<i>herrichi</i>	1906	(Bryk)	C 1
<i>fermata</i>		(Bryk)	=	<i>antoniae</i>	1924	Stauder	C 1
<i>flavibdomaculatus</i>	1921	(Stauder)	=	<i>ladogensis</i>	1911	(Bryk)	B 1
<i>hexamacula</i>	1924	(Stauder)	=	<i>basipunctata</i>	1921	(Belling)	C 1
<i>haltens</i>	1912	(Bryk)	=	<i>cohaerens</i>	1905	(Schultz)	C 1
<i>hyalina</i>	1915	(Bryk)	=	<i>diaphana</i>	1911	(Turati)	D
<i>hypermelas</i>	1913	(Stauder)	=	<i>nox</i>	1921	(Bryk)	C 1
<i>heseboloides</i>	1914	(Schwawerda)	=	<i>immaculata</i>	1913	(Ruhmann)	C 2
<i>inversus</i>	1917	(Kammel)	=	<i>inversa</i>	1898	(Austaut)	E
<i>interruptus</i>	1922	(Lutkemeyer)	=	<i>limbovariegata</i>	1918	(Turati)	C 1
<i>lucundula</i>	1924	(Stauder)	=	<i>intertexta</i>	1906	(Stichel)	B 1
<i>lunigera</i>	1911	(Turati)	=	<i>dentata</i>	1915	(Bryk)	C 2
<i>nonhardi</i>	1912	(Bryk)	=	<i>caeca</i>	1911	(Turati)	B 1
<i>margopupillata</i>	1915	(Bryk)	=	<i>marginepupillata</i>	1918	(Bryk)	A 2
<i>mmemosymoides</i>	1918	(Turati)	=	<i>quincunx</i>	1911	(Bryk)	C 1
<i>monopupillata</i>	1914	(Bryk)	=	<i>unipupillata</i>	1920	(Rebel)	A 1
<i>nigrodivisa</i>	1920	(Riemel)	=	<i>bachmetjevi</i>	1914	(Ugrjumow)	B 2
<i>ochreomaculata</i>	1929	(Bolow)	=	<i>xanthosticta</i>	1924	(Stauder)	B 1
<i>omega-clausum</i>	1924	(Stauder)	=	<i>orbifer</i>	1920	(Rebel)	C 1
<i>perfumigata</i>	1924	(Stauder)	=	<i>fumata</i>	1914	(Vorbrudt)	D
<i>pura</i>	1911	(Turati)	=	<i>defasciata</i>	1913	(Schwawerda)	C 1
<i>posticelongatus</i>	1917	(Kammel)	=	<i>appendiculata</i>	1918	(Turati)	B 2
<i>reductus</i>	1917	(Kammel)	=	<i>expupillata</i>	1911	(Rocci)	B 1
<i>rubromaculata</i>	1929	(Bolow)	=	<i>expupillata</i>	1911	(Rocci)	B 1
<i>rubromaculatus</i>	1917	(Kammel)	=	<i>expupillata</i>	1911	(Rocci)	B 1
<i>septemmacula</i>		(Bryk)	=	<i>heptamacula</i>	1924	(Stauder)	C 1
<i>smidti</i>	1914	(Bryk)	=	<i>expupillata</i>	1911	(Rocci)	B 1
<i>smidtiana</i>	1922	(Bryk)	=	<i>ponsoni</i>	1906	(Culot)	C 1
<i>teniata</i>	1925	(Osthelder)	=	<i>escudei</i>	1923	(Bryk)	C 2
<i>wiskotti</i> (ssp)		(Oberthur)	=	<i>dilatata</i>	1910	Thierry Mieg	E

*Concordance entre les "noms collectifs" d'EISNER et BROWN, et les appellations classiques :*

<i>alboocellata</i>	<i>f. albomaculata</i>	(Muschamp)
<i>analis conjuncta</i>	<i>f. ampliusmaculata</i>	(Verity)
<i>discocircumcincta</i>	<i>f. escudei</i>	(Bryk)
<i>exanalis</i>	<i>f. immaculata</i>	(Ruhmann)
<i>excincta</i>	<i>f. philippsi</i>	(Schutz)
<i>ochreoocellata</i>	<i>f. ochreomaculata</i>	(Kammel)
	= <i>xanthosticta</i>	(Stauder)
<i>rubroocellata</i>	<i>f. rubromaculata</i>	(Kammel)
	= <i>expupillata</i>	(Rocci)
<i>biexcelsior</i>	<i>f. biornata</i>	(Osthelder)